

AD-A164 074

THE STUDY OF SOME EARLY TRANSITION METALS IN OXIDATION  
STATE IV USING THE. (U) TEXAS SOUTHERN UNIV HOUSTON  
B L WILSON 11 OCT 85 ARO-17759.1-CH-H DARG29-88-G-0006

1/1

UNCLASSIFIED

F/G 7/4


NL

END

FILED

JAN

1986



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

2

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD-A164 074

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
ARO 17759.1-CH-H	N/A	N/A
4. TITLE (and Subtitle) The Study of Some Early Transition Metals in Oxidation State IV Using the Potassium Poly(pyrazolyl)borates as Ligands		5. TYPE OF REPORT & PERIOD COVERED 22 Sep 80 - 16 Sep 85 Final Report
6. AUTHOR(s) Bobby L. Wilson		7. PERFORMING ORG. REPORT NUMBER
8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-G-0006		9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
10. PERFORMING ORGANIZATION NAME AND ADDRESS Texas Southern University Houston, TX 77004		11. REPORT DATE October 11, 1985
12. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		13. NUMBER OF PAGES 5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Transition Metals Potassium Poly(Pyrazolyl)Borate Titanium Poly(Pyrazolyl)Borate Zirconium Poly(Pyrazolyl)Borate Hafnium Poly(Pyrazolyl)Borate		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The primary purpose of this investigation was to gain a better understanding of the fundamental chemistry of the early transition elements in their high oxidation states. The work involved the preparation and characterization of titanium, zirconium and hafnium poly(prazolyl)borate complexes with emphasis on the structure, mode of bonding and magnetic behavior of these complexes.		

DTIC  
ELECTE  
FEB 12 1986  
S B D

DTIC FILE COPY

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ARO 17759.1-CH-H

THE STUDY OF SOME EARLY TRANSITION METALS IN  
OXIDATION STATE IV USING THE POTASSIUM  
POLY(PYRAZOLYL)BORATES AS LIGANDS

FINAL REPORT

BOBBY L. WILSON

OCTOBER 11, 1985

U.S. ARMY RESEARCH OFFICE

GRANT NUMBER DAAG-29-80-G-0006

TEXAS SOUTHERN UNIVERSITY  
3100 CLEBURNE AVENUE  
HOUSTON, TEXAS 77004

APPROVED FOR PUBLIC RELEASE:  
DISTRIBUTION UNLIMITED

86 2 7 198

## STATEMENT OF PROBLEM


The primary purpose of this investigation was to gain a better understanding of the fundamental chemistry of the early transition elements in their high oxidation states. The work involved the preparation and characterization of titanium, zirconium and hafnium poly(prazoly1)borate complexes with emphasis on the structure, mode of bonding and magnetic behavior of these complexes.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Special
A-1	



# TETRAKIS(PYRAZOLYL)BORATE COMPLEXES


The reactions of zirconium tetrahalides,  $ZrCl_4$ ,  $ZrBr_4$  and  $ZrI_4$ , and hafnium tetrahalides,  $HfCl_4$  and  $HfI_4$ , with stoichiometric amounts of potassium tetrakis-(pyrazolyl)borate,  $K(B(Pz)_4)$  in dichloromethane or toluene proceeded according to equation I.



The products were isolated as white powders which were quite soluble in dichloromethane and toluene.

Proton magnetic resonance spectra of the complexes were recorded at ambient temperature using  $CD_2Cl_2$  as solvent and  $SiMe_4$  as an internal standard. The spectrum of each complex exhibited three resonances (singlet, doublet and triplet). The observation of only three resonances is an indication that the fully substituted complexes are stereochemically nonrigid. This fluxional behavior was attributed to a tumbling or exchange process.

When two moles of the potassium tetrakis(pyrazolyl)borate were allowed to react with one mole of  $ZrI_4$ ,  $HfCl_4$  or  $HfI_4$  the reactions occurred according to equation II.




Products were isolated as brownish-red, light orange and yellow crystalline solids for the  $ZrI_4$ ,  $HfCl_4$  and  $HfI_4$  respectively. The complexes were air and water sensitive as indicated by color changes when exposed to the atmosphere or placed in water.

The  $^1H$  nmr and  $^{13}C$  nmr spectrum of the zirconium complex each consisted of six and seven resonances respectively that implied a tridentate nature of the  $B(Pz)_4^-$  ligand and therefore eight coordination around zirconium. The spectrum of each hafnium complex exhibited only three resonances. The observation of only three resonances corresponded to four equivalent pyrazolyl groups which

was attributed to a tumbling or exchange process.


### HYDROTRIS(PYRAZOLYL)BORATE COMPLEXES

The reactions of zirconium tetrahalides (Cl and Br) with 1:2 molar amounts of potassium hydrotris(pyrazolyl)borate,  $K(HB(Pz)_3)$ , in dichloromethane proceeded according to the equation III.




The  $ZrCl_2(HB(Pz)_3)_2$  complex was isolated as flaky white crystals which were extremely soluble in dichloromethane.  $ZrCl_2(HB(Pz)_3)_2$  also showed good solubility in chloroform, dimethylformamide, dimethylsulfoxide and was insoluble in benzene and tetrahydrofuran.  $ZrBr_2(HB(Pz)_3)_2$  was isolated as flaky crystals with a pinkish tint. The infrared spectra of both the chloride and bromide complexes were virtually identical with the B-H and aromatic C-H bonds at  $2500\text{ cm}^{-1}$  and  $3110\text{ cm}^{-1}$  respectively. The  $^1H$  nmr spectra were obtained using deuterated chloroform as solvent. The presence of two sets of triplets and doublets for each complex suggested a bidentate nature of the ligand with the formation of eight-coordinate complexes.

White crystalline dichlorobis (hydrotris(pyrazolyl)borate) hafnium (IV) was obtained by the reaction of a 1:2 molar ratio of hafnium tetrachloride with potassium hydrotris(pyrazolyl)borate. The reaction proceeded according to equation IV.



An eight-coordinate structure was assigned based on the  $^1H$  nmr and the infrared spectra.

Tetrakis (hydrotris(pyrazolyl)borate) hafnium (IV) was prepared according to reaction V.



The complex was recovered as a white crystalline powder with a melting point range

of 58° to 60°C. The <sup>1</sup>H nmr spectrum showed only a triplet at δ6.6 and a doublet at δ8.0. The observation of only two peaks implied spectroscopic identify of the three pyrazolyl gfoups and a tumbling process was suggested to explain this spectroscopic identity.

Samples of these and other compounds are available for testing of catalytic characteristics in advanced propellant application.

THE VIEW, OPINIONS, AND/OR FINDINGS CONTAINED IN THIS REPORT ARE THOSE OF THE AUTHOR(S) AND SHOULD NOT BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION, POLICY, OR DECISION, UNLESS SO DESIGNATED BY OTHER DOCUMENTATION.

SCIENTIFIC PERSONEL

Dwight Reid	(M.S.)	1981
Renard L. Thomas	(M.S.)	1981
Renita Y. Webster	(B.S.)	1981
Herbert O. Akalonu	(M.S.)	1982
Elloree Lawson	(B.S.)	1982
Francis I. Addo	(M.S.)	1983
Veronica A. Jefferson	(B.S.)	1984
Samuel A. Ogunkeye	(M.S.)	1984
Charles Asowata	(M.S.)	1985

**END**

**FILMED**

**3-86**

**DTIC**